

# POSITIVELY AWARE 2012 HIV DRUG CHART



Each drug is recommended for use in combination with other antiretrovirals (except possibly **Atripla, Complera, and the "Quad"**). Dose may need to be adjusted; see drug page or package insert for complete information. Drugs shown are actual size, except for Fuzeon. Thanks to Renata Smith, PharmD, University of Illinois at Chicago. Drug chart information is **current as of Feb. 1, 2012**. For a list of Walgreens pharmacies that specialize in HIV/AIDS care and other resources, visit [www.HIV.walgreens.com/chart](http://www.HIV.walgreens.com/chart). You can also scan the QR code at right with your iPhone or other smartphone to go directly to the website. You can also click on a drug's name or its picture to go to that drug's page in the HIV Drug Guide. See the POSITIVELY AWARE 2012 HIV Drug Guide at [www.positivelyaware.com/drugguide](http://www.positivelyaware.com/drugguide) for more information.



SPONSORED BY  
**Walgreens**

NUCLEOSIDE/NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITOR (NRTI, NRTTI)



## Combivir

lamivudine / zidovudine, or **3TC / AZT**

**STANDARD DOSE:** One tablet (150 mg lamivudine / 300 mg zidovudine), twice daily. No food restrictions (with or without food).



## EpiVir

lamivudine, or **3TC**

**STANDARD DOSE:** One 300 mg tablet, once daily; or one 150 mg tablet twice daily. No food restrictions (with or without food).



## Retrovir

zidovudine, **AZT**, or **ZDV**

**STANDARD DOSE:** One 300 mg tablet, twice daily. No food restrictions (with or without food).



## Truvada

emtricitabine / tenofovir, or **FTC / TDF**

**STANDARD DOSE:** One tablet (200 mg emtricitabine / 300 mg tenofovir), once daily. No food restrictions (with or without food).



## Viread

tenofovir disoproxil fumarate, or **TDF**

**STANDARD DOSE:** One 300 mg tablet, once daily. No food restrictions (with or without food).



## Ziagen

abacavir sulfate, or **ABC**

**STANDARD DOSE:** Two 300 mg tablets, once daily (or one 300 mg tablet twice daily). No food restrictions (with or without food).



## Intelence

etravirine, or **ETV**

**STANDARD DOSE:** One 200 mg tablet, twice daily, or two 100 mg tablets, twice daily. Take with food.



## Sustiva

efavirenz, or **EFV**

**STANDARD DOSE:** One 600 mg tablet, once daily or three 200 mg capsules, once daily. Take on an empty stomach, or may be taken with a light, low-fat snack.



## Aptivus

tipranavir, or **TPV**

**STANDARD DOSE:** Two 250 mg capsules with two 100 mg tablets Norvir, twice daily. Aptivus must be taken with Norvir. Take with food with Norvir tablets. No food restrictions with Norvir capsules, but take with food to improve Norvir tolerability.



## Invirase

saquinavir, or **SQV**

**STANDARD DOSE:** Two 500 mg tablets with one 100 mg tablet Norvir, twice daily. Invirase must be taken with Norvir. Take with food, or within two hours after a meal.



## Lexiva

fosamprenavir calcium, or **FPV**

**STANDARD DOSE:** Two 700 mg tablets plus 100 mg or 200 mg Norvir, once daily or two 700 mg tablets, twice daily (if protease inhibitor experienced, 700 mg with 100 mg Norvir, twice daily). No food restrictions (with or without food).



## Prezista

darunavir, or **DRV**

**STANDARD DOSE:** Two 400 mg tablets, once a day for first-time therapy, or one 600 mg tablet, twice a day for treatment-experienced people. All doses taken with 100 mg Norvir. Take with food.



## Viracept

nelfinavir, or **NFV**

**STANDARD DOSE:** Two 625 mg tablets, twice daily (or five 250 mg tablets, twice daily). Take with a meal.

## dolutegravir

dolutegravir, or **DTG**

**STANDARD DOSE:** One 50 mg tablet once daily chosen for Phase 3 research. Twice-daily dosing was shown to work better in people who have resistance to Isentress and the experimental elvitegravir in a short study, and a large Phase 3 trial will look at the twice-daily dose. Expected to have no food restrictions. (Investigational drug at press time; photo unavailable.)



## Fuzeon

enfuvirtide, **T-20**, or **ENF**

**STANDARD DOSE:** 90 mg, twice daily subcutaneous injections. No food restrictions (with or without food).



## Atripla

efavirenz / emtricitabine / tenofovir, or **EFV / FTC / TDF**

**STANDARD DOSE:** One tablet (600 mg efavirenz / 200 mg emtricitabine / 300 mg tenofovir), once daily. Take on an empty stomach, or may be taken with a light, low-fat snack.

## The "Quad"

elvitegravir / cobicistat / emtricitabine / tenofovir, or **EVG / COBI / FTC / TDF**

**STANDARD DOSE:** One tablet (150 mg elvitegravir / 150 mg cobicistat / 200 mg emtricitabine / 300 mg tenofovir) once daily has been selected in Phase 3 studies. Take with food. (Not yet approved at press time; photo unavailable.)

## Complera

rilpivirine / emtricitabine / tenofovir, or **RPV / FTC / TDF**

**STANDARD DOSE:** One tablet (25 mg rilpivirine / 200 mg emtricitabine / 300 mg tenofovir) once daily. Take with food (with a meal of at least 400 calories).

## cobicistat

(formerly **GS-9350**, or **COBI**)

**STANDARD DOSE:** Not yet established. 150 mg used in research. Cobicistat is not an HIV drug, but is used to increase (boost) the levels of elvitegravir (an investigational integrase inhibitor, see elvitegravir) and HIV protease inhibitors. (Not yet approved at press time; photo unavailable.)

NUCLEOSIDE/NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITOR (NRTI, NRTTI)

## Emtriva

emtricitabine, or **FTC**

**STANDARD DOSE:** One 200 mg capsule, once daily. No food restrictions (with or without food).

## Epzicom

abacavir / lamivudine, or **ABC / 3TC**

**STANDARD DOSE:** One tablet (600 mg abacavir / 300 mg lamivudine), once daily. No food restrictions (with or without food).

## Trizivir

abacavir / lamivudine / zidovudine, or **ABC / 3TC / AZT**

**STANDARD DOSE:** One tablet (300 mg abacavir / 150 mg lamivudine / 300 mg zidovudine), twice daily. No food restrictions (with or without food).

## Videx EC

didanosine, or **ddl**

**STANDARD DOSE:** One 400 mg capsule, once daily with adjustments for weight and drug interactions (see the Drug Guide). Take on an empty stomach, one hour before or two hours after food or drink, except water. (Generic pictured.)

## Zerit

stavudine, or **d4T**

**STANDARD DOSE:** One 40 mg capsule, twice daily with adjustments for weight (see the Drug Guide). No food restrictions (with or without food).

## Edurant

rilpivirine hydrochloride, or **RPV**

**STANDARD DOSE:** One 25 mg tablet, once daily. Take with food (with a meal of at least 400 calories).

## Rescriptor

delavirdine, or **DLV**

**STANDARD DOSE:** Two 200 mg tablets, three times a day (or four 100 mg tablets, three times a day). No food restrictions (can be taken with or without food).

## Viramune XR

nevirapine, or **NVP**

**STANDARD DOSE:** One 200 mg IR (immediate release) tablet daily for the first 14 days, then one 400 mg tablet once daily for Viramune XR, or one 200 mg IR tablet twice daily, or two 200 mg IR tablets once daily (once daily not FDA approved for IR formulation). No food restrictions (with or without food).

## Crixivan

indinavir, or **IDV**

**STANDARD DOSE:** Two 400 mg capsules, every 8 hours, or two 400 mg capsules with 100 mg Norvir twice a day. Take with no food or a low-fat snack (with food with Norvir); drink at least 1.5 liters of water daily.

## Kaletra

lopinavir / ritonavir, or **LPV / r**

**STANDARD DOSE:** Four tablets (200 mg lopinavir / 50 mg ritonavir), once daily if fewer than 3 lopinavir resistance mutations, or two 200 / 50 mg tablets, twice daily. Half-strength film-coated tablet available: 100 mg lopinavir / 25 mg ritonavir. No food restrictions; take with food to improve Norvir tolerability.

## Norvir

ritonavir, or **RTV**

**STANDARD DOSE:** Used mostly for boosting, 100 mg–400 mg dosed once or twice a day with another PI. Must be taken with food.

## Reyataz

atazanavir sulfate, or **ATV**

**STANDARD DOSE:** One 300 mg capsule plus 100 mg Norvir, once daily or two 200 mg capsules (without Norvir), once daily for treatment-naïve adults (see Drug Guide). Take with food.

## Isentress

raltegravir, or **RAL**

**STANDARD DOSE:** One 400 mg tablet, twice daily. No food restrictions (with or without food).

## elvitegravir

elvitegravir, or **EVG**

**STANDARD DOSE:** One 150 mg or one 85 mg tablet once a day boosted with either Norvir or cobicistat. The dose depends on drug interactions with the co-administered PI. Take with food. (Not yet approved at press time; photo unavailable.)

## Selzentry

maraviroc, or **MVC**

**STANDARD DOSE:** 150, 300, or 600 mg (available in 150 and 300 mg tablets), twice daily, depends on other medications used (see Drug Guide). No food restrictions (with or without food).

## Complera

rilpivirine / emtricitabine / tenofovir, or **RPV / FTC / TDF**

**STANDARD DOSE:** One tablet (25 mg rilpivirine / 200 mg emtricitabine / 300 mg tenofovir) once daily. Take with food (with a meal of at least 400 calories).

NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITOR (NNRTI)

NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITOR (NNRTI)

PROTEASE INHIBITOR (PI)

PROTEASE INHIBITOR (PI)

INTEGRASE INHIBITOR

INTEGRASE INHIBITOR

ENTRY INHIBITOR

ENTRY INHIBITOR

SINGLE-TABLET REGIMEN

SINGLE-TABLET REGIMEN

PK ENHANCER

PK ENHANCER